



in form on the farm®

# PREMIN® ANTISTERIL 2010

Good reproduction parameters and health calves rearing are two basic presumptions for effective dairy cattle breeding. Therefore, top dairy farmers apply Premin® Antisteril two weeks before calving and during the first third of lactation. It contains minerals, microelements, vitamins and  $\beta$ -caroten.

## IMPORTANCE:

- specially formulated vitamin-mineral feed to improve reproduction parameters
- decreases post-partum disorders
- minimises afterbirth placenta retention
- supports physiologic uterus involution
- improves ovarian function and heat induction
- reduces appearance of ovary atrophy and ovarian cyst syndrom
- supports immune system of the animal
- reduces appearance of ketosis and endrometris
- supplements highly bioavailable microelements
- has antioxidant effect
- contains CLA,  $\beta$ -caroten, vitamins A, D3 and E

*Better reproduction parameters*



VVS Verměřovice s.r.o.  
Krmivářská 225  
561 52 Verměřovice

+420 465 642 670  
GSM: +420 775 755 175  
email: vvs@vvs.cz, [www.vvs.cz](http://www.vvs.cz)



## APPLICATION INDICATIONS:

- poor reproduction parameters
- problems with urogenital system
- metabolic disorders

## COMPOSITION:

vitamins A, D3 and E, niacin,  $\beta$ -caroten, organic Selenium, organic microelements, macroelements, CLA

## DOSAGE:

50 - 100 g per head per day

## RECOMMENDATION:

- in a dry and transition period – 3 weeks before and 3-4 weeks after calving
- during the first third of lactation
- apply to all cows for 1-2 months when health problems are indicated



VVS Verměřovice s.r.o.  
Krmivářská 225  
561 52 Verměřovice

+420 465 642 670  
GSM: +420 775 755 175  
email: vvs@vvs.cz, [www.vvs.cz](http://www.vvs.cz)

